

Procedure Instruction Packet

READ this instruction packet completely at least 7 days prior to your procedure!!

**Your procedure may be cancelled if these instructions are not followed.

Date of Procedure: _____

Procedure Scheduled: _____

Required Arrival Time: _____

> We ask all patients to arrive 1 hour prior to the estimated start time of their procedure.

Estimated Start Time of Procedure: _____

Note: The exact time for each procedure varies from patient to patient. Our physicians give each patient outstanding care and undivided attention.

Please print this packet out if you are receiving it by email.

It is important that you read the instructions a few times before your procedure. We recommend you set a reminder on your calendar 7-10 days prior to your procedure to ensure you are prepared and understand your restrictions.

Important: If you are scheduled for a colonoscopy, *follow our instructions* <u>ONLY.</u> Do NOT follow the instructions within the colonoscopy prep kit you receive at the pharmacy.

Following the instructions within this packet is very important. We want to ensure your procedure will be performed with success.

You will receive a confirmation call or text 72 hours prior to your scheduled procedure.

Colonoscopy Prep Instructions Colyte Split Dose Prescription

10 Days Prior	1 Week Prior	5 Days Prior	2 Days Prior	1 Day Prior	Procedure Day
You must	Pick up your		Drink at least four 8oz.	Begin clear liquid. <u>NO Solid foods</u> today. Red,	Morning Dose Colyte- Part 2
STOP taking	prescription.	<u>Stop</u>	glasses of water	purple or blue colored liquids are not	
Phentermine, or any medication containing Phentermine. <u>Examples:</u> Qsymia, Qnexa, Adipex-P, Suprenza, Fastin or Phentercot.	Only follow these instructions. DoNOT follow box instructions.If you take dietary, herbal or fiber supplements, or medications containing iron, discontinue these 7 days before your appointment.If you take Coumadin, Plavix, Pradaxa, Xarelto, Eliquis or other	consuming all food with seeds, corn and nuts. <u>Stop</u> taking ibuprofen, Advil, Aleve and NSAIDs You may take Tylenol as needed.	Advil, NSAIDsBiod Pressure and Heart medications as taking NSAIDsAcceptable clear liquids include: water, clear broth, apple juice, white cranberry juice, white grape juice, soda, gelatin (Jello), popsicles, coffee, and tea (no milk or creamer).If your pro At 3am Dr until rest or litishStop taking directed. Take them taking ourorfen, Advil, NSAIDsBlood Pressure and Heart medications as directed. Take them the day of your procedure with a less than 2 hours before procedure.If you are taking diabetic medication, cut your dose in half this day and do not take any diabetic medication day of procedure. If you have an insulin pump, check with prescribing MD for instructions.If your pro At 3am Dr until rest or It show the grape juice, soda, gelatin (Jello), popsicles, coffee, and tea (no milk or creamer).If your pro At 730am minutes up consumed lise than 2 hours before procedure.You may takeInhalers and Nasal Spray: Continue to take theseDrink at least four 8oz. glasses of water throughout the day. Do NOT drink alcohol the	If your procedure is scheduled before 1130am: At 3am Drink 8oz of solution every 15 minutes until rest of solution has been consumed. It should take 1 hour to complete If your procedure is scheduled after 1130am: At 730am Drink 8oz of solution every 15 minutes until rest of solution every 15 minutes until rest of solution has been consumed. It should take 1 hour to complete **You need to finish 4 hours prior to your procedure. Nothing by mouth at least 4 hours prior to scheduled procedure time-including gum, hard candy or mints.	
	blood thinners: You must obtain permission to withhold this medication prior to your procedure. Contact our pre- operative nurse for concerns or questions at 410 224-3636 You MAY continue to take: once daily aspirin (81mg or 325mg daily)		procedure. If you have a fever, cold, or respiratory symptoms within 48 hours of your procedure- call us. Eat light dinner. NO Solid Foods after midnight	 5pm: Begin Colyte Prep- Part 1 Step 1- Add cool water to line on container. Close and shake to dissolve. Add Flavor Pack to Container if not already added. Step 2- Drink 8oz of solution every 15 minutes until 75% (3/4) has been consumed. You should be done by 9pm. Step 3- Store the remainder of the solution in the refrigerator overnight. 	You may take your essential morning medications with a small sip of water, <u>at</u> <u>least</u> 2 hours prior to your procedure, unless otherwise directed by your physician. <u>You will need:</u> -Insurance Cards -Driver's License/Photo ID -Any co-insurance fees due -A responsible adult driver to drive you home. A taxi or shuttle is not an approved means of transportation <u>unless you have</u> a family member or friend with you. You may not drive until the day after your procedure.

YOUR BOWEL Prep IS VERY IMPORTANT- Please read instructions carefully!

Medication Record Form

I	1