

YOUR COLONOSCOPY IS SCHEDULED FOR:

e of Procedure:	
stro Health Physician:	
ival Time:	
ility/address:	
one:	
gistration Assistant:	

Appointments missed or canceled with less than 72 hours' notice may result in a fee in accordance with applicable office policies.

SAME DAY-CLEAR LIQUID PREPARATION INSTRUCTIONS FOR YOUR **COLONOSCOPY WITH PLENVU**

PLEASE REVIEW THE INSTRUCTIONS AT LEAST 3 DAYS PRIOR TO THE PROCEDURE USE THESE PREPARATION INSTRUCTIONS AND NOT THOSE INCLUDED WITH THE PLENVU BOX.



It is important that you follow these directions carefully and complete the entire preparation to ensure the most effective cleansing.

- If you are taking any blood thinners and/or insulin AND this was NOT discussed with your doctor at the time of scheduling the procedure, please call us **BEFORE** starting the preparation. These include, but are not limited to Aspirin, Coumadin-(Warfarin), Plavix-(Clopidogrel), Xarelto-(Rivaroxaban), Pradaxa-(Dabigatran), Elequis-(Apixaban), Arixtra-(Fondaparinux), Effient-(Prasugrel), Aggrenox-(Aspirin/Dipirydamole), and all NSAIDs (Ibuprofen, Motrin, Aleve, Naproxen, Celebrex, Mobic).
- YOU MUST CONTINUE TO TAKE ALL your other medications every day prior to the colonoscopy AND take them with a small sip of water on the day of your colonoscopy.
- **IF YOU ARE DIABETIC, DO NOT** take your diabetic pills the evening prior or day of the colonoscopy.
- Please plan to have someone drive you home after the procedure. You will not be allowed to drive yourself home.
- You cannot take an Uber, Lyft, taxi, or public transportation home from your procedure without an adult companion.

1.855.GASTRO1 GastroHealth.com



SAME DAY PREPARATION INSTRUCTIONS FOR YOUR COLONOSCOPY WITH PLENVU



3 DAYS BEFORE THE PROCEDURE

STOP eating fruits and vegetables including beans, corn and seeds.

THE DAY PRIOR TO THE PROCEDURE

You may not have any solid food. If desired you may have a liquid yogurt only for breakfast before 10 AM, no later. After 10 am you may have only clear liquids. For example: water, coffee, tea, broth, sodas, apple juice, Gatorade, Kool-aid, Jell-O, popsicles, and hard candy.

DO NOT DRINK MILK. DO NOT DRINK LIQUIDS THAT ARE RED OR PURPLE IN COLOR.

THE MORNING OF THE PROCEDURE

STEP 1:

For best results mix ahead and refrigerate the solution prior to drinking.

- Empty **Dose 1** into the mixing container that comes with your PLENVU.
- Add lukewarm water (16 oz) to the top line of the container and mix or shake to completely dissolve.



STEP 2:


_____, (7 hours before arrival time) begin drinking Dose 1. Finish the entire contents within 30 minutes.

STEP 3:

- Rinse the mixing container with water. Refill to the fill line (16 oz) with any clear liquid of your choice.
- Drink the entire contents of the mixing container over the next 30 minutes.
- Continue drinking additional clear liquids. (At least 16 ounces)

STEP 4:

Follow same directions as Step 1 using **Dose 2** pouches A and B.



STEP 5:

(4 hours before the arrival time) begin drinking Dose 2. Finish the entire contents Starting at ___ within 30 minutes.

STEP 6:

- Rinse the mixing container with water. Refill to the fill line (16 oz) with any clear liquid of your choice.
- Drink the entire contents of the mixing container over the next 30 minutes.
- Continue drinking additional clear liquids. (At least 16 ounces).

YOU MUST COMPLETE STEP 6 AND STOP ALL LIQUIDS BY (At least 2 hours before your Arrival Time) Please call us if you have any questions or if you are having difficulty with the preparation.

> 1.855.GASTRO1 GastroHealth.com