

#### YOUR COLONOSCOPY IS SCHEDULED FOR:

e of Procedure:	
tro Health Physician:	
val Time:	
ility/address:	
ne:	
sistration Assistant:	

# DAY BEFORE-CLEAR LIQUID PREPARATION INSTRUCTIONS FOR YOUR COLONOSCOPY WITH CLENPIQ.

PLEASE REVIEW THE INSTRUCTIONS *AT LEAST 3 DAYS PRIOR* TO THE PROCEDURE USE THESE PREPARATION INSTRUCTIONS *AND NOT* THOSE INCLUDED WITH CLENPIQ.



### It is important that you follow these directions carefully and complete the entire preparation to ensure the most effective cleansing.

- If you are taking any blood thinners and/or insulin AND this was NOT discussed with your doctor at the time of scheduling the procedure, please call us BEFORE starting the preparation. These include, but are not limited to Aspirin, Coumadin-(Warfarin), Plavix-(Clopidogrel), Xarelto-(Rivaroxaban), Arixtra-(Fondaparinux), Effient-(Prasugrel), Aggrenox-(Aspirin/Dipirydamole), and all NSAIDs (Ibuprofen, Motrin, Aleve, Naproxen, Celebrex, Mobic).
- **YOU MUST CONTINUE TO TAKE ALL** your other medications every day prior to the colonoscopy AND take them with a small sip of water on the day of your colonoscopy.
- IF YOU ARE DIABETIC, DO NOT take your diabetic pills the evening prior or day of the colonoscopy.
- Please plan to have someone drive you home after the procedure. You will not be allowed to drive yourself home.
- You cannot take an Uber, Lyft or taxi home from your procedure without an adult companion.





## DAY BEFORE PREPARATION INSTRUCTIONS FOR YOUR COLONOSCOPY WITH CLENPIQ



#### **3 DAYS BEFORE THE PROCEDURE**

STOP eating fruits and vegetables including beans, corn and seeds.

#### THE DAY PRIOR TO THE PROCEDURE

You may not have any solid food. If desired, you may have a liquid yogurt only for breakfast before 10 AM, no later. After 10 am, you may have only clear liquids. For example: water, coffee, tea, broth, sodas, apple juice, Gatorade, Kool-aid, Jell-O, popsicles, and hard candy. It is important to drink extra clear liquids before your prep to maintain adequate hydration.

**BEGIN YOUR CLENPIQ** 

#### DO NOT DRINK MILK. DO NOT DRINK LIQUIDS THAT ARE RED OR PURPLE IN COLOR.

#### Step 1:

Starting between 5:00 PM and 7:00 PM, drink ONE (1) bottle of Clenpiq. Drink entire content from the bottle.

#### Step 2:

Follow with FIVE (5) 8-ounce glasses of any clear liquid. About one (1) cup every 30 minutes.

# 8 oz 8 oz 8 oz 8 oz

#### THE DAY OF THE PROCEDURE

#### Step 3:

Starting at \_\_\_\_\_\_ (4 hours before the arrival time) drink ONE (1) bottle of Clenpiq. Drink entire content from the bottle.

#### Step 4:

Follow with THREE (3) 8-ounce glasses of any clear liquid. About 1 cup every 3 minutes. Finish the entire prep even if stool is clear liquid.



100 Plost Corn Lete Stel Tallo Stot All Ligotos Di (At icast 2 nouis belote your Allivat init	YOU MUST COMPLETE STEP 4 AND STOP ALL LIQUIDS BY (	At least 2 hours b	efore you	ur Arrival Time
-----------------------------------------------------------------------------------------------	----------------------------------------------------	--------------------	-----------	-----------------

Please call us if you have any questions or if you are having difficulty with the preparation. We are available 24 hours a day, 7 days a week.

