

YOUR COLONOSCOPY IS SCHEDULED FOR:

Date of Procedure:
Gastro Health Physician:
Arrival Time:
acility/address:
Phone:
Registration Assistant:

SAME DAY-CLEAR LIQUID PREPARATION INSTRUCTIONS FOR YOUR COLONOSCOPY WITH CLENPIQ.

PLEASE REVIEW THE INSTRUCTIONS **AT LEAST 3 DAYS PRIOR** TO THE PROCEDURE. USE THESE PREPARATION INSTRUCTIONS **AND NOT** THOSE INCLUDED WITH THE CLENPIQ BOX.



It is important that you follow these directions carefully and complete the entire preparation to ensure the most effective cleansing.

- If you are taking any blood thinners **AND** this was **NOT** discussed with your doctor at the time of scheduling the procedure, please call us **BEFORE** starting the preparation. These include, but are not limited to Aspirin, Coumadin-(Warfarin), Plavix-(Clopidogrel), Xarelto-(Rivaroxaban), Arixtra-(Fondaparinux), E˜ent-(P rasugrel), Aggrenox-(Aspirin/Dipirydamole), and all NSAIDs (for example Ibuprofen, Motrin, Aleve, Naproxen, Celebrex, Mobic).
- YOU MUST CONTINUE TO TAKE ALL your other medications every day prior to the colonoscopy AND take them with a small sip of water on the day of your colonoscopy.
- IF YOU ARE DIABETIC, DO NOT take your diabetic pills the day of the colonoscopy.
- Please make arrangements to have someone drive you home after the procedure. You will not be allowed to drive yourself home.





SAME DAY PREPARATION INSTRUCTIONS FOR YOUR COLONOSCOPY WITH CLENPIQ.



3 DAYS BEFORE THE PROCEDURE

STOP eating fruits and vegetables including beans, corn and seeds.

THE DAY PRIOR TO THE PROCEDURE

You may not have any solid food. If desired you may have a liquid yourt only for breakfast before 10 AM, no later. After 10 am you may have only clear liquids. For example: water, coffee, tea, broth, sodas, apple juice, Gatorade, Kool-aid, Jell-O, popsicles, and hard candy.

It is important to drink extra clear liquids before your prep to maintain adequate hydration. DO NOT DRINK MILK. DO NOT DRINK LIQUIDS THAT ARE RED OR PURPLE IN COLOR.

THE MORNING OF THE PROCEDURE BEGIN YOUR CLENPIQ

STEP 1:

Starting at

(7 hours before the arrival time)

- Drink one (1) bottle of Clenpiq.
- Drink the entire content right from the bottle.

STEP 2:

- Follow with FIVE (5) 8-ounce glasses of any clear liquid.
- About 1 cup every 30 minutes.



STEP 3:

Starting at:

(4 hours before arrival time)

- Drink the second bottle of Clenpiq
- Drink the entire content right from the bottle.

STEP 4:

- Follow with THREE (3) 8-ounce cups of any clear liquid.
- About 1 cup every 30 minutes.
- Finish the entire prep even if stool is clear liquid.



YOU MUST COMPLETE STEP 4 AND STOP ALL LIQUIDS BY

(At least 2 hours before your Arrival Time)

Please call us if you have any questions or if you are having difficulty with the preparation. We are available 24 hours a day, 7 days a week.

